

4. With respect to a fixed origin  $O$  the lines  $l_1$  and  $l_2$  are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 11 \\ 2 \\ 17 \end{pmatrix} + \lambda \begin{pmatrix} -2 \\ 1 \\ -4 \end{pmatrix} \quad l_2: \mathbf{r} = \begin{pmatrix} -5 \\ 11 \\ p \end{pmatrix} + \mu \begin{pmatrix} q \\ 2 \\ 2 \end{pmatrix}$$

Jan  
2009

where  $\lambda$  and  $\mu$  are parameters and  $p$  and  $q$  are constants. Given that  $l_1$  and  $l_2$  are perpendicular,

- (a) show that  $q = -3$ .

(2)

Given further that  $l_1$  and  $l_2$  intersect, find

- (b) the value of  $p$ ,

(6)

- (c) the coordinates of the point of intersection.

(2)

The point  $A$  lies on  $l_1$  and has position vector  $\begin{pmatrix} 9 \\ 3 \\ 13 \end{pmatrix}$ . The point  $C$  lies on  $l_2$ .

Given that a circle, with centre  $C$ , cuts the line  $l_1$  at the points  $A$  and  $B$ ,

- (d) find the position vector of  $B$ .

(3)

---

4. The line  $l_1$  has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix}$$

and the line  $l_2$  has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -4 \\ 1 \end{pmatrix}$$

where  $\lambda$  and  $\mu$  are parameters.

The lines  $l_1$  and  $l_2$  intersect at the point  $A$  and the acute angle between  $l_1$  and  $l_2$  is  $\theta$ .

- (a) Write down the coordinates of  $A$ .

(1)

- (b) Find the value of  $\cos \theta$ .

(3)

The point  $X$  lies on  $l_1$  where  $\lambda = 4$ .

- (c) Find the coordinates of  $X$ .

(1)

- (d) Find the vector  $\overrightarrow{AX}$ .

(2)

- (e) Hence, or otherwise, show that  $|\overrightarrow{AX}| = 4\sqrt{26}$ .

(2)

The point  $Y$  lies on  $l_2$ . Given that the vector  $\overrightarrow{YX}$  is perpendicular to  $l_1$ ,

- (f) find the length of  $AY$ , giving your answer to 3 significant figures.

(3)

Jan  
2011

4. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$  and the point  $B$  has position vector  $-2\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ . The points  $A$  and  $B$  lie on a straight line  $l$ .

(a) Find  $\overrightarrow{AB}$ . (2)

(b) Find a vector equation of  $l$ . (2)

The point  $C$  has position vector  $2\mathbf{i} + p\mathbf{j} - 4\mathbf{k}$  with respect to  $O$ , where  $p$  is a constant.

Given that  $AC$  is perpendicular to  $l$ , find

(c) the value of  $p$ , (4)

(d) the distance  $AC$ . (2)

---

7. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $(2\mathbf{i} - \mathbf{j} + 5\mathbf{k})$ , the point  $B$  has position vector  $(5\mathbf{i} + 2\mathbf{j} + 10\mathbf{k})$ , and the point  $D$  has position vector  $(-\mathbf{i} + \mathbf{j} + 4\mathbf{k})$ .

The line  $l$  passes through the points  $A$  and  $B$ .

- (a) Find the vector  $\overrightarrow{AB}$ . (2)
- (b) Find a vector equation for the line  $l$ . (2)
- (c) Show that the size of the angle  $BAD$  is  $109^\circ$ , to the nearest degree. (4)

The points  $A$ ,  $B$  and  $D$ , together with a point  $C$ , are the vertices of the parallelogram  $ABCD$ , where  $\overrightarrow{AB} = \overrightarrow{DC}$ .

- (d) Find the position vector of  $C$ . (2)
- (e) Find the area of the parallelogram  $ABCD$ , giving your answer to 3 significant figures. (3)
- (f) Find the shortest distance from the point  $D$  to the line  $l$ , giving your answer to 3 significant figures. (2)



6. With respect to a fixed origin  $O$ , the lines  $l_1$  and  $l_2$  are given by the equations

$$l_1 : \mathbf{r} = (-9\mathbf{i} + 10\mathbf{k}) + \lambda(2\mathbf{i} + \mathbf{j} - \mathbf{k})$$

$$l_2 : \mathbf{r} = (3\mathbf{i} + \mathbf{j} + 17\mathbf{k}) + \mu(3\mathbf{i} - \mathbf{j} + 5\mathbf{k})$$

where  $\lambda$  and  $\mu$  are scalar parameters.

- (a) Show that  $l_1$  and  $l_2$  meet and find the position vector of their point of intersection. (6)

- (b) Show that  $l_1$  and  $l_2$  are perpendicular to each other. (2)

The point  $A$  has position vector  $5\mathbf{i} + 7\mathbf{j} + 3\mathbf{k}$ .

- (c) Show that  $A$  lies on  $l_1$ . (1)

The point  $B$  is the image of  $A$  after reflection in the line  $l_2$ .

- (d) Find the position vector of  $B$ . (3)
-

June  
2009

7. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $(8\mathbf{i} + 13\mathbf{j} - 2\mathbf{k})$ , the point  $B$  has position vector  $(10\mathbf{i} + 14\mathbf{j} - 4\mathbf{k})$ , and the point  $C$  has position vector  $(9\mathbf{i} + 9\mathbf{j} + 6\mathbf{k})$ .

The line  $l$  passes through the points  $A$  and  $B$ .

- (a) Find a vector equation for the line  $l$ . (3)

- (b) Find  $|\overrightarrow{CB}|$ . (2)

- (c) Find the size of the acute angle between the line segment  $CB$  and the line  $l$ , giving your answer in degrees to 1 decimal place. (3)

- (d) Find the shortest distance from the point  $C$  to the line  $l$ . (3)

The point  $X$  lies on  $l$ . Given that the vector  $\overrightarrow{CX}$  is perpendicular to  $l$ ,

- (e) find the area of the triangle  $CXB$ , giving your answer to 3 significant figures. (3)
-

June  
2010

7. The line  $l_1$  has equation  $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ , where  $\lambda$  is a scalar parameter.

The line  $l_2$  has equation  $\mathbf{r} = \begin{pmatrix} 0 \\ 9 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}$ , where  $\mu$  is a scalar parameter.

Given that  $l_1$  and  $l_2$  meet at the point  $C$ , find

(a) the coordinates of  $C$ .

(3)

The point  $A$  is the point on  $l_1$  where  $\lambda = 0$  and the point  $B$  is the point on  $l_2$  where  $\mu = -1$ .

(b) Find the size of the angle  $ACB$ . Give your answer in degrees to 2 decimal places.

(4)

(c) Hence, or otherwise, find the area of the triangle  $ABC$ .

(5)

---

6. With respect to a fixed origin  $O$ , the lines  $l_1$  and  $l_2$  are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + \lambda \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} -5 \\ 15 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix},$$

where  $\lambda$  and  $\mu$  are scalar parameters.

(a) Show that  $l_1$  and  $l_2$  meet and find the position vector of their point of intersection  $A$ . (6)

(b) Find, to the nearest  $0.1^\circ$ , the acute angle between  $l_1$  and  $l_2$ . (3)

The point  $B$  has position vector  $\begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix}$ .

(c) Show that  $B$  lies on  $l_1$ . (1)

(d) Find the shortest distance from  $B$  to the line  $l_2$ , giving your answer to 3 significant figures. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

